

# Exact Solution of a Many-Fermion System and Its Associated Boson Field

Cite as: Journal of Mathematical Physics **6**, 304 (1965); <https://doi.org/10.1063/1.1704281>  
Submitted: 22 September 1964 . Published Online: 22 December 2004

Daniel C. Mattis, and Elliott H. Lieb



View Online



Export Citation

## ARTICLES YOU MAY BE INTERESTED IN

### [An Exactly Soluble Model of a Many-Fermion System](#)

Journal of Mathematical Physics **4**, 1154 (1963); <https://doi.org/10.1063/1.1704046>

### [Relationship between Systems of Impenetrable Bosons and Fermions in One Dimension](#)

Journal of Mathematical Physics **1**, 516 (1960); <https://doi.org/10.1063/1.1703687>

### [Periodic table for topological insulators and superconductors](#)

AIP Conference Proceedings **1134**, 22 (2009); <https://doi.org/10.1063/1.3149495>



Journal of  
Mathematical Physics

READ TODAY!

Special Issue: XIXth International  
Congress on Mathematical Physics

# Exact Solution of a Many-Fermion System and Its Associated Boson Field

DANIEL C. MATTIS

*International Business Machines Corp., Thomas J. Watson Research Center,  
Yorktown Heights, New York*

AND

ELLIOTT H. LIEB\*

*Belfer Graduate School of Science, Yeshiva University, New York, New York*

(Received 22 September 1964)

Luttinger's exactly soluble model of a one-dimensional many-fermion system is discussed. We show that he did not solve his model properly because of the paradoxical fact that the density operator commutators  $[\rho(p), \rho(-p')]$ , which always vanish for any finite number of particles, no longer vanish in the field-theoretic limit of a filled Dirac sea. In fact the operators  $\rho(p)$  define a boson field which is *ipso facto* associated with the Fermi-Dirac field. We then use this observation to solve the model, and obtain the exact (and now nontrivial) spectrum, free energy, and dielectric constant. This we also extend to more realistic interactions in an Appendix. We calculate the Fermi surface parameter  $\bar{n}_k$ , and find:  $\partial \bar{n}_k / \partial k|_{k_F} = \infty$  (i.e., there exists a sharp Fermi surface) only in the case of a sufficiently weak interaction.

## I. INTRODUCTION

THE search for a soluble but realistic model in the many-electron problem has been just about as unfruitful as the historic quest for the philosopher's stone, but has equally resulted in valuable byproducts. For example, 15 years ago Tomonaga<sup>1</sup> published a theory of interacting fermions which was soluble only in one dimension with the provision that certain truncations and approximations were introduced into his operators. Nevertheless he had success in showing approximate boson-like behavior of certain collective excitations, which he identified as "phonons." (Today we would denote these as "plasmons," following the work of Bohm and Pines.<sup>2</sup>) Lately, Luttinger<sup>3</sup> has revived interest in the subject by publishing a variant model of spinless and massless one-dimensional interacting fermions, which demonstrated a singularity at the Fermi surface, compatible with the results of the modern many-body perturbation theory.<sup>4</sup>

Unfortunately, in calculating the energies and wavefunctions of his model Hamiltonian, Luttinger fell prey to a subtle paradox inherent in quantum field theory<sup>5</sup> and therefore did not achieve a correct

*solution of the problem he himself had posed.* In the present paper we shall give the solution to his interesting problem and calculate the free energy. We shall show the existence of collective plasmon modes, and shall calculate the singularity at the Fermi surface (which may in fact disappear if the interaction is strong enough), the energy of the plasmons, and the (nontrivial) dielectric constant of the system. In an Appendix we shall show how the model may be generalized in such a manner as to remove certain restrictions on the interactions which Luttinger had found necessary to impose.

It is fortunate that solid-state and many-body theorists have so far been spared the plagues of quantum field theory. Second quantization has been often just a convenient bookkeeping arrangement to save us from writing out large determinantal wavefunctions. However there is a difference between very large determinants and *infinitely* large ones; we shall show that one of the important differences is the failure of certain commutators to vanish in the field-theoretic limit when common sense and experience based on finite  $N$  tells us they *should* vanish! (Here  $N$  refers to the number of particles in the field.)

\* Research supported by the U. S. Air Force Office of Scientific Research.

<sup>1</sup> S. Tomonaga, *Progr. Theoret. Phys. (Kyoto)* **5**, 544 (1950).

<sup>2</sup> D. Bohm and D. Pines, *Phys. Rev.* **92**, 609 (1953).

<sup>3</sup> J. M. Luttinger, *J. Math. Phys.* **4**, 1154 (1963). Note that we set his  $v_0 = 1$ , thereby fixing the unit of energy. References to this paper will be frequent, and will be denoted by L (72), for example, signifying his Eq. (72).

<sup>4</sup> J. M. Luttinger and J. C. Ward, *Phys. Rev.* **118**, 1417 (1960).

<sup>5</sup> Luttinger made a transformation, L (8), which was canonical in appearance only. But in the language of G. Barton [*Introduction to Advanced Field Theory*, (Interscience

Publishers, Inc., New York, 1963), pp. 126 *et seq.*] this transformation connected two "unitarily inequivalent" Hilbert spaces, which has as a consequence that commutators, among other operators, must be reworked so as to be well-ordered in fermion field operators. It was first observed by Julian Schwinger [*Phys. Rev. Letters* **3**, 296 (1959)] that the very fact that one postulates the existence of a ground state (i.e., the filled Fermi sea) forces certain commutators to be nonvanishing even though in first quantization they automatically vanish. The "paradoxical contradictions" of which Schwinger speaks seem to anticipate the difficulties in the Luttinger model.

We shall show that these nonvanishing commutators *define* boson fields which must *ipso facto* always be associated with a Fermi-Dirac field, and we shall use the ensuing commutation relations to solve Luttinger's model exactly. Because this model is soluble both in the Hilbert space of finite  $N$  and also in the Hilbert space  $N = \infty$ , with different physical behavior in each, we believe it has applications to the *theory of fields* which go beyond the study of the many-electron problem. The model can be extended to the case of electrons with spin. This has interesting consequences in the band *theory of ferromagnetism*, as will be discussed in some detail in an article under preparation.<sup>5a</sup>

## II. MODEL HAMILTONIAN

We recall Luttinger's Hamiltonian<sup>3</sup> and recapitulate some of his results:

$$H = H_0 + H', \quad (2.1)$$

where the "unperturbed" part is

$$H_0 = \int_0^L dx \psi^\dagger(x) \sigma_3 p \psi(x) \quad (2.2a)$$

$$= \sum_k (a_{1k}^* a_{1k} - a_{2k}^* a_{2k}) k, \quad (2.2b)$$

and the interaction is

$$H' = 2\lambda \iint_0^L dx dy \psi_1^\dagger(x) \psi_1(x) \times V(x-y) \psi_2^\dagger(y) \psi_2(y) \quad (2.3a)$$

$$= \frac{2\lambda}{L} \sum \delta_{k_1+k_2, k_3+k_4} v(k_3 - k_4) \times a_{1k_1}^* a_{1k_2} a_{2k_3}^* a_{2k_4}. \quad (2.3b)$$

Here  $\psi$  is a two-component field and the form (b) of the operator is obtained from (a) by setting

$$\psi = \frac{1}{\sqrt{L}} \sum_k e^{ikz} \begin{pmatrix} a_{1k} \\ a_{2k} \end{pmatrix}$$

and

$$\psi^\dagger = \frac{1}{\sqrt{L}} \sum_k e^{-ikz} (a_{1k}^*, a_{2k}^*), \quad (2.4)$$

with  $a_{ik}$ 's defined to be anticommuting fermion operators which obey the usual relations

$$a_{ik} a_{i'k'} + a_{i'k'} a_{ik} \equiv \{a_{ik}, a_{i'k'}\} = 0 \quad (2.5)$$

$$\{a_{i',k}^*, a_{j',k'}^*\} = 0, \text{ and } \{a_{ik}, a_{j',k'}^*\} = \delta_{ij'} \delta_{kk'}.$$

Luttinger noted that for an appropriate operator

$S_0$ , the canonical transformation

$$\tilde{H} = e^{i\lambda S_0} H e^{-i\lambda S_0}. \quad (2.6)$$

gave the result that

$$\tilde{H} = H_0, \quad (2.7)$$

and consequently that *the spectrum of  $H = H_0 + H'$  was the same as that of  $H_0$ , independent of the interaction  $V(x-y)$* . This can be explicitly verified for his choice of

$$S_0 = \iint_0^L dx dy \psi_1^\dagger(x) \psi_1(x) E(x-y) \psi_2^\dagger(y) \psi_2(y), \quad (2.8)$$

where  $E(x)$ , not to be confused with the energy  $E$ , is defined by:

$$\partial E(x-y)/\partial x \equiv V(x-y), \quad (2.9)$$

assuming that

$$\bar{V} \equiv \frac{1}{L} \int_0^L V(x) dx = 0. \quad (2.10)$$

In the Appendix we shall show among other things how to generalize to  $\bar{V} \neq 0$ . It is also simple and instructive to verify Eqs. (2.6) and (2.7) somewhat differently by using the *first* quantization,

$$H_0 = -i \sum_{n=1}^N \frac{\partial}{\partial x_n} + i \sum_{m=1}^M \frac{\partial}{\partial y_m} \quad (2.11)$$

and

$$H' = 2\lambda \sum_{n=1}^N \sum_{m=1}^M V(x_n - y_m), \quad (2.12)$$

where  $N$  and  $M$  are, respectively, the total number of "1" particles and "2" particles, with coordinates  $x_n$  and  $y_m$ , respectively. The properly antisymmetrized wavefunctions are given by

$$\Psi = \det |e^{ik_i x_i}| \det |e^{i a_j y_j}| \times \exp \left\{ \sum_{n=1}^N \sum_{m=1}^M i E(x_n - y_m) \right\}. \quad (2.13)$$

Using Eqs. (2.9) and (2.10),  $\Psi$  is readily seen to obey Schrödinger's equation

$$H\Psi = E\Psi \quad (2.14)$$

with just the unperturbed eigenvalue

$$E = \sum_{n=1}^N k_n - \sum_{m=1}^M q_m. \quad (2.15)$$

The wavenumbers are of the form

$$k_i \text{ or } q_j = 2\pi \text{ integer}/L, \quad (2.16)$$

as required for periodic boundary conditions. This is in exact agreement with the results of Ref. 3, and can also be checked in perturbation theory; first-

<sup>5a</sup> D. Mattis, *Physics* 1, 184 (1964).

order perturbation theory also gives vanishing results, and indeed, it is easy to verify that to every order in  $\lambda$  the cancellation is complete, in accordance with the exact result given above.

Up to this point, Luttinger's analysis (which we have briefly summarized) is perfectly correct. It is the next step that leads to difficulty. The Hamiltonian discussed so far has no ground-state energy; in order to remove this obstacle, and thereby establish contact with a real electron gas, Luttinger proposed modifying the model by "filling the infinite sea" of negative energy levels (i.e., all states with  $k_1 < 0$  and  $q_2 > 0$ ). Following L(8) we define  $b$ 's and  $c$ 's obeying the usual anticommutators, such that

$$\text{and } \begin{aligned} a_{1k} &= \begin{cases} b_k & k \geq 0 \\ c_k^* & k < 0, \end{cases} \\ a_{2k} &= \begin{cases} b_k & k < 0 \\ c_k^* & k \geq 0. \end{cases} \end{aligned} \quad (2.17)$$

Using this notation the total particle-number operator becomes

$$\mathcal{N} = \sum_{\text{all } k} b_k^* b_k - c_k^* c_k \quad (2.17a)$$

(i.e., the number of particles minus the number of holes).

Since the Hamiltonian commutes with  $\mathcal{N}$  we can demand that  $\mathcal{N}$  have eigenvalue  $N_0$ . In the noninteracting ground state there are no holes and the  $b$  particles are filled from  $-k_F$  to  $k_F$  where  $k_F = \pi(N_0/L) = \pi\rho$ . The noninteracting ground-state energy is  $N_0\pi\rho + \text{energy of the filled sea } (W)$ .

The kinetic energy assumes the form

$$H_0 = \sum_{\text{all } k} (b_k^* b_k + c_k^* c_k) |k| + W, \quad (2.18)$$

where

$$W = \left( \sum_{k < 0} k - \sum_{k > 0} k \right) \quad (2.18a)$$

is the infinite energy of the filled sea, an uninteresting  $c$  number which we drop henceforth in accordance with Luttinger's prescription. The interaction  $[H', \text{Eq. (2.3)}]$  and the operator  $S_0$ , Eq. (2.8)] can also be expressed in the new language by means of the substitution (2.17). The reader will no doubt be surprised, as indeed we were, to find that now with the new operators, Eq. (2.7), with  $\tilde{H}$  defined in (2.6), is no longer obeyed.

Upon further reflection one sees that this must be so, on the basis of very general arguments. In the new Hilbert space defined by the transformation to the particle-hole language (2.17),  $H$  is no longer unbounded from below and now has a ground state.

A general and inescapable *concavity theorem* states that if  $E_0(\lambda)$  is the ground-state energy in the presence of interactions, (2.3), then

$$\partial^2 E_0(\lambda) / \partial \lambda^2 < 0. \quad (2.19)$$

This inequality is incompatible with the previous result, viz. all  $E =$  independent of  $\lambda$ , which was possible only in the strange case of a system without a ground state.

The same thing can be seen more trivially using second-order perturbation theory (first-order perturbation theory vanishes). It is easily seen that

$$E_0^{(2)} = - \left( \frac{2\lambda}{L} \right)^2 \sum_k \frac{|v(k)|^2}{2k} n_1(k) n_2(-k), \quad (2.20)$$

where  $n_1(k)$  and  $n_2(k)$  are the number of ways of shifting a particle of type "1" and type "2" respectively by an amount  $k$  to an unoccupied state. A simple geometric exercise will convince the reader of the following facts: (1) if we start with a state having a finite number of particles, then  $n_1$  and  $n_2$  are *always* even functions of  $k$  (i.e., there are just as many ways to increase the momentum by  $k$  as to decrease it by the same amount.) (2) If we start with a filled infinite sea then there is no way to decrease the momentum of the "1" particles nor to increase the momentum of "2" particles. Hence for this second case  $n_1(k)n_2(-k)$  is nonzero only for  $k > 0$ . Thus  $E_0^{(2)}$  vanishes for a state with a finite number of particles, but it is negative for a filled sea.

If the reader is unconvinced by perturbation theory, then he can easily prove that  $E_0$  is lowered by doing a variational calculation.

What has gone wrong? We turn to some algebra to resolve this paradox, and following this, present a solution of the field-theoretic problem defined by  $H_0 + H'$  in the representation of  $b$ 's and  $c$ 's.

### III. CASE OF THE FILLED DIRAC SEA

The various relevant operators are given below; the form (a) of each equation will *not* be used in the bulk of the paper, and is just given here for completeness. In the following equations,  $p > 0$ .

$$\rho_1(+p) \equiv \sum_k a_{1, k+p}^* a_{1, k} \quad (3.1a)$$

$$= \sum_{k < -p} c_{k+p} c_k^* + \sum_{-p \leq k < 0} b_{k+p}^* c_k^* + \sum_{k \geq 0} b_{k+p}^* b_k, \quad (3.1b)$$

$$\rho_1(-p) \equiv \sum_k a_{1, k}^* a_{1, k+p} \quad (3.2a)$$

$$= \sum_{k < -p} c_k c_{k+p}^* + \sum_{-p \leq k < 0} c_k b_{k+p} + \sum_{k \geq 0} b_k^* b_{k+p}, \quad (3.2b)$$

$$\rho_2(+p) \equiv \sum_k a_{2k+p}^* a_{2k} \quad (3.3a)$$

$$= \sum_{k < -p} b_{k+p}^* b_k + \sum_{-p \leq k < 0} c_{k+p} b_k + \sum_{k \geq 0} a_{k+p} c_k^* \quad (3.3b)$$

$$\rho_2(-p) \equiv \sum_k a_{2k}^* a_{2k+p} \quad (3.4a)$$

$$= \sum_{k < -p} b_k^* b_{k+p} + \sum_{-p \leq k < 0} b_k^* c_{k+p}^* + \sum_{k > 0} c_k c_{k+p}^* \quad (3.4b)$$

Equations (3.1a)–(3.4a) give the density operators in the original representation, so let us calculate in this language a commutator such as (assume  $p \geq p' \geq 0$  for definiteness)

$$\begin{aligned} [\rho_1(-p), \rho_1(p')] &= \sum_{k, k'} [a_{1k}^* a_{1k+p}, a_{1k'+p'}^* a_{1k'}] \\ &= \sum_{k=-\infty}^{+\infty} a_{1k}^* a_{1k+p-p'} - \sum_{k=-\infty}^{+\infty} a_{1k+p'}^* a_{1k+p} = 0. \end{aligned} \quad (3.5)$$

The zero result could have been expected by writing the operators in first quantization:

$$\rho_1(-p) = \sum_n e^{-ipx_n} \quad \text{and} \quad \rho_2(p) = \sum_m e^{ipx_m}, \quad (3.6)$$

whence they evidently commute. Nevertheless, the zero result is achieved in (3.5) only through the almost ‘‘accidental’’ cancellation of two operators, each of which may diverge in the field-theory limit when  $N = \infty$ . We now show that in that limit the operators in fact no longer cancel, by evaluating the commutator using form (b) for the density operators. It is a matter of only some minor manipulation to obtain the important new result:

$$\begin{aligned} [\rho_1(-p), \rho_1(p')] &= [\rho_2(p), \rho_2(-p')] \\ &= \delta_{p, p'} \sum_{-p < k < 0} 1 = \frac{pL}{2\pi} \delta_{p, p'}, \quad (p' > 0). \end{aligned} \quad (3.7a)$$

In addition,

$$[\rho_1(p), \rho_2(p')] = 0. \quad (3.7b)$$

A quick check is provided by evaluating the vacuum expectation value

$$\begin{aligned} \langle 0 | [\rho_1(-p), \rho_1(p)] | 0 \rangle \\ = \sum_{-p < k, k' < 0} \langle 0 | c_k b_{k+p} b_{k'}^* c_{k'+p}^* | 0 \rangle = pL/2\pi, \end{aligned} \quad (3.8)$$

which is exactly what is expected on the basis of the previous equation. Evidently the form (b) of the operators  $(2\pi/pL)^{\dagger} \rho_1(+p)$  and  $(2\pi/pL)^{\dagger} \rho_2(-p)$  have properties of boson raising operators [call them  $A^*(p)$  and  $B^*(-p)$ ] and  $(2\pi/pL)^{\dagger} \rho_1(-p)$  and  $(2\pi/pL)^{\dagger} \rho_2(+p)$  have properties of boson lowering operators [ $A(p)$  and  $B(-p)$ ], i.e.,

$$[A, B] = [A^*, B] = 0, \quad (3.9)$$

$$[A(p), A^*(p')] = [B(-p), B^*(-p)] = \delta_{p, p'}.$$

The  $B$  field is the continuation of the  $A$  field to negative  $p$ ; therefore together they form a *single* boson field defined for all  $p$ .

The relationship of the  $\rho(p)$ 's to Luttinger's  $N(x)$ 's, L(25), is obtained by using (2.4):

$$N_1(x) = \psi_1^\dagger(x) \psi_1(x) = \frac{1}{L} \sum_p \rho_1(p) e^{-ipx}, \quad (3.10)$$

$$N_2(x) = \psi_2^\dagger(x) \psi_2(x) = \frac{1}{L} \sum_p \rho_2(p) e^{-ipx}.$$

#### IV. SOLUTIONS OF THE MODEL HAMILTONIAN

Before making use of the results of the previous section, we remark that  $\rho_1(+p)$  and  $\rho_2(-p)$  are exact raising operators of  $H_0$ , and  $\rho_1(-p)$  and  $\rho_2(p)$  are exact lowering operators of  $H_0$  corresponding to excitation energies  $p$ . That is,

$$[H_0, \rho_1(\pm p)] = \pm p \rho_1(\pm p), \quad (4.1)$$

$$[H_0, \rho_2(\pm p)] = \mp p \rho_2(\pm p).$$

The identification of the  $\rho$ 's with boson operators made in the previous section suggested to us the possibility of constructing a new operator  $T$  which obeys the same equations (4.1), as  $H_0$ . This is indeed possible, if we define  $T$  as follows:

$$T \equiv \frac{2\pi}{L} \sum_{p>0} \{ \rho_1(p) \rho_1(-p) + \rho_2(-p) \rho_2(p) \} \quad (4.2)$$

[the  $\rho$ 's being defined here and in the remainder of the paper by Eqs. (3.1b)–(3.4b), i.e., always in the hole-particle representation]. It follows that

$$[T, \rho_1(\pm p)] = \pm p \rho_1(\pm p) \quad (4.3)$$

as required, and similarly for  $\rho_2(\mp p)$ . Therefore, let us decompose  $H$  into two parts

$$H = H_1 + H_2 \quad (4.4)$$

with

$$\begin{aligned} H_1 = H_0 - T &= \left\{ \sum_k |k| (b_k^* b_k + c_k^* c_k) \right. \\ &\quad \left. - \frac{2\pi}{L} \sum_{p>0} \{ \rho_1(p) \rho_1(-p) + \rho_2(-p) \rho_2(p) \} \right\}, \end{aligned} \quad (4.5)$$

and

$$\begin{aligned} H_2 = H' + T \\ = \frac{1}{L} \left[ 2\lambda \sum_{p>0} \{ v(p) \rho_1(-p) \rho_2(p) + v(-p) \rho_1(p) \rho_2(-p) \} \right. \\ \left. + 2\pi \sum_{p>0} \{ \rho_1(p) \rho_1(-p) + \rho_2(-p) \rho_2(p) \} \right] \end{aligned} \quad (4.6)$$

with  $v(p) = \text{real}$ , even function of  $p$ . By actual construction, all the  $\rho$  operators which appear in  $H_2$

commute with  $H_1$ . This will be an important feature in constructing an exact solution of the model. We define an Hermitian operator  $S$ ,

$$S = \frac{2\pi i}{L} \sum_{\alpha, \beta} \frac{\varphi(p)}{p} \rho_1(p) \rho_2(-p), \quad (4.7)$$

where  $\varphi(p)$  is also a real, even, function of  $p$  to be determined subsequently by imposing a condition that the unitary transformation  $e^{iS}$  diagonalize  $H_2$ . First we evaluate the effect of such a transformation on various operators. It commutes with  $H_1$ ,

$$e^{iS} H_1 e^{-iS} = H_1 = H_0 - T, \quad (4.8)$$

because both  $\rho_1$  and  $\rho_2$  appearing in  $S$  commute with  $H_1$ , as noted above. In the following,  $p$  can have either sign:

$$e^{iS} \rho_1(p) e^{-iS} = \rho_1(p) \cosh \varphi(p) + \rho_2(p) \sinh \varphi(p), \quad (4.9)$$

$$e^{iS} \rho_2(p) e^{-iS} = \rho_2(p) \cosh \varphi(p) + \rho_1(p) \sinh \varphi(p). \quad (4.10)$$

We have verified that this transformation is a proper unitary transformation and preserves commutation relations (3.7) as well as anticommutation relations (2.5), and the reader may easily check this point.  $H_2$  is brought into canonical form by requiring that in  $(\exp iS) H_2 (\exp -iS)$  there be no cross terms such as  $\rho_1(p) \rho_2(-p)$ . This leads to the equation

$$\tanh 2\varphi = -\lambda v(p)/\pi, \quad (4.11)$$

which cannot be obeyed unless

$$|\lambda v(p)| < \pi \quad \text{for all } p. \quad (4.12)$$

Equation (4.12) serves to limit the magnitude of potentials capable of having well-behaved solutions (e.g., a real ground-state energy). For the more realistic potentials discussed in the Appendix, there is also a more realistic bound on  $v(p)$ : there,  $v(p)$  may not be *too* attractive, but it can have any magnitude when it is repulsive, i.e., positive.

With the choice of  $\varphi$  in (4.11), the evaluation of  $H_2$  becomes

$$e^{iS} H_2 e^{-iS} = \frac{2\pi}{L} \sum_{p>0} \text{sech } 2\varphi(p) \{ \rho_1(p) \rho_1(-p) + \rho_2(-p) \rho_2(p) \} - \sum_{p>0} p(1 - \text{sech } 2\varphi). \quad (4.13a)$$

The second term is the vacuum renormalization energy

$$W_1 = - \sum_{p>0} p(1 - \text{sech } 2\varphi) = \frac{L}{2\pi} \int_0^\infty dp p \left\{ \left( 1 - \frac{\lambda^2 v^2(p)}{\pi^2} \right)^{\frac{1}{2}} - 1 \right\}. \quad (4.13b)$$

It may be expanded in powers of  $\lambda$  to effect a comparison with Goldstone's many-body perturbation theory<sup>4</sup>; we have checked that they agree to third order.

The problem is now formally solved, for we can find all the eigenfunctions and eigenvalues by studying Eqs. (4.4), (4.8), and (4.13). First notice that the operator  $T$  does not depend upon the interaction and that if there is *no interaction* we could write the Hamiltonian either as

$$H = H_0, \quad (4.14a)$$

or as

$$H = (H_0 - T) + T = H_1 + H_2. \quad (4.14b)$$

Since  $H_1$  and  $H_2$  commute, every eigenstate,  $\Psi$ , of  $H$  may be assumed to be an eigenfunction of  $H_1$  and  $H_2$  separately. Moreover,  $\Psi$  may also be assumed to be an eigenfunction of each  $\alpha_p = A_p^+ A_p$  and  $\beta_p = B_{-p}^+ B_{-p}$  for all  $p > 0$ , since these operators commute with  $H$  and  $\mathcal{N}$ .

Evidently (4.14a) and (4.14b) provide two different ways of viewing the noninteracting spectrum.  $H_0$  is quite degenerate: the raising operators of  $H_0$  are the  $b^+$ 's and  $c^+$ 's. By requiring that  $\Psi$  also be an eigenstate of  $\alpha_p$ ,  $\beta_p$  and  $H$ , we are merely attaching quantum numbers to the degenerate levels of  $H_0$ . If  $\alpha_p \Psi = n_p \Psi$  and  $\beta_p \Psi = m_p \Psi$  (where  $n_p$  and  $m_p$  are of course integers), we say that we have  $n_p$  plasmons of momentum  $p$  and  $m_p$  plasmons of momentum  $-p$ . With no interaction the energy of a plasmon is

$$\epsilon(p) = |p|. \quad (4.15)$$

We may speak of  $H_1$  as the quasiparticle part of the Hamiltonian; in  $H_1$  the operator  $T$  plays the role of subtracting the plasmon part of the energy from  $H_0$ .

When we turn on the interaction, the above description of the energy levels is still valid, except that now we are *forced* to use the form (4.14b) because  $H_2$  is no longer  $T$ . The degeneracy of  $H$  is partially removed by the interaction, because now the energy of a plasmon is

$$\epsilon'(p) = |p| \text{sech } 2\varphi(p). \quad (4.16)$$

Notice that the plasmon energy is always *lowered* [and therefore the plasmons cannot propagate faster than the speed of light  $c = 1$ , i.e.,  $d\epsilon'/dp \leq 1$ . In the more realistic case discussed in the Appendix, the plasmon energy *can* be increased by the interaction although  $d\epsilon'/dp \leq 1$  is always obeyed.] by the interaction; if (4.12) is violated the plasmon energy is no longer real and the system becomes unstable. Note, there are no plasmons in the ground state, so that  $W_1$  (4.13), is the shift in the ground-state energy of the system.

There is one important point, however, that requires some elucidation. We would like to be able to say that in view of the fact that  $H_1$ ,  $\alpha(p)$ , and  $\beta(p)$

conserve particle number, the most general energy level of  $H$  (fixed  $N_0$ ) is the sum of *any* energy of  $H_1$  (same  $N_0$ , and no plasmons) plus *any* (plasmon) energy of  $H_2$  (note: the plasmon spectrum is independent of  $N_0$ ). Were we dealing with a finite-dimensional vector space, such a statement would not be true, for even though  $H_1$  and  $H_2$  commute they could not possibly be independent. Thus, if  $H_2$  had  $n$  eigenvalues  $e_1, \dots, e_n$ , and if  $H_1$  had an equal number  $E_1, \dots, E_n$  the general total eigenvalue would not be *any* combination of  $e_i + E_i$  for this would give too many values (viz.  $n^2$  instead of  $n$ .) But we are dealing with an infinite-dimensional Hilbert space and the additivity hypothesis is in fact true for the present model.

To prove this assertion we consider any eigenstate  $\Psi$  which is necessarily parameterized by the integers  $n_p$  and  $m_p$ . Consider the state  $\Phi = \{\prod_p (A_p)^{n_p} (B_p)^{m_p}\} \Psi$ . The state  $\Phi$  is nonvanishing and has quantum numbers  $n_p = 0 = m_p$ . It is also an eigenstate of  $H_1$  with energy  $E_1(\Psi)$ . In addition (and this is the important point) the state  $\Psi$  may be recovered from  $\Phi$  by the equation

$$\Psi = \text{const} \times \left\{ \prod_p (A_p^+)^{n_p} (B_p^+)^{m_p} \right\} \Phi.$$

To every state  $\Psi$ , therefore, there corresponds a *unique* state  $\Phi$  from which it may be obtained using raising operators. Conversely, to any eigenstate of  $H_1$  (for fixed  $N_0$ ) we may apply raising operators as often as we please and obtain a new (nonvanishing) eigenstate. Thus the general energy is an arbitrary sum of quasiparticle and plasmon energies.

It may be wondered where we used the fact that the Hilbert space is infinite-dimensional in the above proof. The answer lies in the boson commutation relations of the  $A$ 's and  $B$ 's. It is impossible to have such relations in a finite-dimensional vector space.

The eigenvalues corresponding to these states  $\Phi$  will be labeled in some order,  $E_i$  ( $i = 1, 2, \dots$ ), so that the total canonical partition function  $Z(\lambda)$  and the free energy  $F(\lambda)$  are given by

$$\begin{aligned} Z(\lambda) &= e^{-F(\lambda)/kT} \\ &= \left( \sum_i e^{-E_i/kT} \right) (e^{-W_1/kT}) \prod_{\substack{\text{all } p \\ p \neq 0}} \left( \sum_{n=0}^{\infty} e^{-n\epsilon(p)/kT} \right). \end{aligned} \quad (4.17)$$

The first factor is difficult to evaluate directly. However it can be obtained circuitously by noting that the energies  $E_i$  are independent of  $\lambda$  and therefore

$$\begin{aligned} Z(0) &= e^{-F(0)/kT} \\ &= \left( \sum_i e^{-E_i/kT} \right) \prod_{\substack{\text{all } p \\ p \neq 0}} \left( \sum_{n=0}^{\infty} e^{-n\epsilon(p)/kT} \right). \end{aligned} \quad (4.18)$$

But the second factor can be trivially evaluated, as can  $F(0) =$  free energy of noninteracting fermions. Therefore we use (4.18) to eliminate the trace involving the  $E_i$ 's in (4.17), with the final result:

$$\begin{aligned} F(\lambda) &= F(0) + W_1 \\ &+ 2kT \sum_{p>0} \ln \left\{ (1 - e^{-\epsilon'(p)/kT}) / (1 - e^{-\epsilon(p)/kT}) \right\}, \end{aligned} \quad (4.19)$$

where  $\epsilon$  and  $\epsilon'$  are given in (4.15) and (4.16). It is noteworthy that the ground state and free energy both diverge in the case of a  $\delta$ -function potential.

## V. EVALUATION OF THE MOMENTUM DISTRIBUTION

In this section we calculate the mean number of particles with momentum  $k$ . This quantity is  $\bar{n}_k$  and is the expectation value of

$$n_k = b_k^\dagger b_k \quad (5.1)$$

in the ground state. Since  $\bar{n}_k$  is an even function of  $k$  we need only consider  $k > 0$ , and it is further convenient to introduce a Fourier transform so that [using (2.4)]

$$\bar{n}_k = \frac{1}{L} \iint_0^L ds dt e^{ik(s-t)} I(s, t). \quad (5.2)$$

Here

$$\begin{aligned} I(s, t) &= \langle \Psi | \psi_1^\dagger(s) \psi_1(t) | \Psi \rangle \\ &= \langle \Psi_0 | e^{iS} \psi_1^\dagger(s) e^{-iS} e^{iS} \psi_1(t) e^{-iS} | \Psi_0 \rangle, \end{aligned} \quad (5.3)$$

where  $S$  is given by (4.7),  $\Psi$  is the new ground state, and  $\Psi_0$  is the noninteracting ground state which is filled with  $b$  particles between  $-k_F$  and  $k_F$  and has no holes (or  $c$  particles). This assignment depends on there having been no level crossing, which can be readily verified using (4.7)-(4.13).

In order to calculate the quantity  $e^{iS} \psi_1(t) e^{-iS}$  we introduce the auxiliary operator

$$f_\sigma(t) = e^{i\sigma S} \psi_1(t) e^{-i\sigma S}, \quad (5.4)$$

where  $\sigma$  is a  $c$  number. We observe that  $f_1(t)$  is the desired quantity while

$$f_0(t) = \psi_1(t). \quad (5.5)$$

In addition,

$$\begin{aligned} \partial f / \partial \sigma &= e^{i\sigma S} i[S, \psi_1(t)] e^{-i\sigma S} \\ &= e^{i\sigma S} [2\pi/L \sum_p \rho_2(-p) \varphi(p) p^{-1} e^{ip t}] e^{-i\sigma S} f_\sigma(t), \end{aligned} \quad (5.6)$$

where we have used the commutation relations (3.7) as well as the fact that  $\psi_1$  commutes with  $\rho_2$ . Equa-

tion (5.6) is a differential equation for  $f_\sigma(t)$  and (5.5) is the boundary condition. The solution is

$$f_\sigma(t) = W_\sigma(t)R_\sigma(t)\psi_1(t), \quad (5.7)$$

where

$$W_\sigma(t) = \exp \left\{ 2\pi/L \sum_{p>0} [\rho_1(-p)e^{ip\sigma t} - \rho_1(p)e^{-ip\sigma t}] p^{-1} [\cosh \sigma\varphi(p) - 1] \right\} \quad (5.8)$$

and

$$R_\sigma(t) = \exp \left\{ 2\pi/L \sum_{p>0} [\rho_2(-p)e^{ip\sigma t} - \rho_2(p)e^{-ip\sigma t}] p^{-1} \sinh \sigma\varphi(p) \right\} \quad (5.9)$$

The reader may verify that (5.7) satisfies (5.5) and (5.6) by using the commutation relations (3.7). We recall the well-known rule that

$$\exp(A+B) = \exp(A)\exp(B)\exp(-1/2[A,B]) \quad (5.10)$$

when  $[A, B]$  commutes with  $A$  and  $B$ . From here on we shall set  $\sigma = 1$  and drop it as a subscript. We note that since  $\rho_1(p)^+ = \rho_1(-p)$  and  $\rho_2(p)^+ = \rho_2(-p)$ ,

$$R^+(t) = R^{-1}(t) \quad \text{and} \quad W^+(t) = W^{-1}(t). \quad (5.11)$$

We also note that  $R$  and  $W$  commute with each other. Thus, (5.3) becomes

$$I(s, t) = \langle \Psi_0 | \psi_1^+(s)R^{-1}(s)W^{-1}(s)W(t)R(t)\psi_1(t) | \Psi_0 \rangle \\ = I_1(s, t)I_2(s, t), \quad (5.12)$$

where

$$I_1(s, t) = \langle \Psi_1 | \psi_1^+(s)W^{-1}(s)W(t)\psi_1(t) | \Psi_1 \rangle, \quad (5.13)$$

$$I_2(s, t) = \langle \Psi_2 | R^{-1}(s)R(t) | \Psi_2 \rangle.$$

We have used the fact that the ground state is a product state:  $\Psi_0 = \Psi_1 * \Psi_2$  where  $\Psi_1$  is a state of the "1" field and  $\Psi_2$  is a state of the "2" field.  $\Psi_1$  is filled with  $b$  particles up to  $+k_F$  and has no  $c$  particles;  $\Psi_2$  is filled with  $b$  particles down to  $-k_F$  and has no  $c$  particles.

Now, using the definition (5.8) and the rule (5.10) we easily find that

$$W^{-1}(s)W(t) = W_-(s, t)W_+(s, t)Z_1(s, t), \quad (5.14)$$

with

$$W_+(s, t) = \exp \left\{ 2\pi/L \sum_{p>0} \rho_1(-p) [\cosh \varphi(p) - 1] \right. \\ \left. \times p^{-1} (e^{ip\sigma t} - e^{ip\sigma s}) \right\},$$

$$W_-(s, t) = \exp \left\{ 2\pi/L \sum_{p>0} \rho_1(p) [\cosh \varphi(p) - 1] \right. \\ \left. \times p^{-1} (e^{-ip\sigma s} - e^{-ip\sigma t}) \right\},$$

$$Z_1(s, t) = \exp \left\{ 2\pi/L \sum_{p>0} [\cosh \varphi(p) - 1]^2 \right. \\ \left. \times p^{-1} (e^{ip\sigma(t-s)} - 1) \right\}. \quad (5.15)$$

Likewise,

$$R^{-1}(s)R(t) = R_-(s, t)R_+(s, t)Z_2(s, t), \quad (5.16)$$

with

$$R_+(s, t) = \exp \left\{ 2\pi/L \sum_{p>0} \rho_2(p) [\sinh \varphi(p)] \right. \\ \left. \times p^{-1} (e^{-ip\sigma s} - e^{-ip\sigma t}) \right\},$$

$$R_-(s, t) = \exp \left\{ 2\pi/L \sum_{p>0} \rho_2(-p) [\sinh \varphi(p)] \right. \\ \left. \times p^{-1} (e^{ip\sigma t} - e^{ip\sigma s}) \right\},$$

$$Z_2(s, t) = \exp \left\{ 2\pi/L \sum_{p>0} [\sinh \varphi(p)]^2 \right. \\ \left. \times p^{-1} (e^{ip\sigma(t-s)} - 1) \right\}. \quad (5.17)$$

We see at once from the definition (3.1b), (3.2b), of  $\rho_1(p)$  that, for  $p > 0$ ,  $\rho_1(-p) |\Psi_1\rangle = 0$ . Similarly  $\langle \Psi_1 | \rho(p) = 0$ ,  $\rho_2(p) |\Psi_2\rangle = 0$ , and  $\langle \Psi_2 | \rho_2(-p) = 0$ . Hence,

$$I_2(s, t) = Z_2(s, t)$$

and

$$I_1(s, t) = Z_1(s, t) \langle \Psi_1 | W^{-1}\psi_1^+(s)W_+W_+\psi_1(t)W_+^{-1} | \Psi_1 \rangle. \quad (5.18)$$

If we now define

$$h_+(y) = 2\pi/L \sum_{p>0} [\cosh \varphi(p) - 1] \\ \times p^{-1} (e^{ip\sigma t} - e^{ip\sigma s}) e^{-ip\sigma y}, \\ h_-(y) = 2\pi/L \sum_{p>0} [\cosh \varphi(p) - 1] \\ \times p^{-1} (e^{-ip\sigma t} - e^{-ip\sigma s}) e^{ip\sigma y}, \quad (5.19)$$

combining (3.10) and (5.15) we have that

$$W_+(s, t) = \exp \int_0^L N_1(y) h_+(y) dy, \quad (5.20)$$

$$W_-(s, t) = \exp - \int_0^L N_1(y) h_-(y) dy.$$

Since

$$[\psi_1(x), N_1(y)] = \delta(x-y)\psi_1(x), \quad (5.21)$$

$$[\psi_1^+(x), N_1(y)] = -\delta(x-y)\psi_1^+(x),$$

it follows that

$$W_+(s, t)\psi_1(t)W_+^{-1}(s, t) = \psi_1(t) \exp [-h_+(t)] \quad (5.22)$$

$$W_+^{-1}(s, t)\psi_1^+(s)W_+(s, t) = \psi_1^+(s) \exp [+h_-(s)].$$



Finally,

$$\begin{aligned} \langle \Psi_1 | \psi_1^\dagger(s) \psi_1(t) | \Psi_1 \rangle &= 1/L \sum_{p \leq k_F} e^{ip(t-s)} \\ &\equiv Z_3(s, t). \end{aligned} \quad (5.23)$$

Combining all these results, we conclude that

$$I(s, t) = Z_0(s, t) Z_1(s, t) Z_2(s, t) Z_3(s, t), \quad (5.24)$$

where

$$\begin{aligned} Z_0(s, t) &= \exp(h_-(s) - h_+(t)) \\ &= \exp \left\{ -4\pi/L \sum_{p>0} [\cosh \varphi(p) - 1] \right. \\ &\quad \left. \times (1 - e^{ip(s-t)}) \right\}. \end{aligned} \quad (5.25)$$

In order to make a comparison with Luttinger's calculation of  $\bar{n}_k$ , we first observe that the functions  $Z_i(s, t)$  are really functions of  $r = s - t$  and that they are periodic in  $s$  and  $t$  in  $(0, L)$ . We then define the functions  $G(r)$  and  $Q(r)$  as follows:

$$\exp[-Q(r)] \equiv G(r) \equiv Z_0(r) Z_1(r) Z_2(r). \quad (5.26)$$

Substituting (5.26), (5.24), and (5.23) into (5.2) we obtain

$$\bar{n}_k = 2\pi/L \sum_{p \leq k_F} F(k - p), \quad (5.27)$$

where

$$F(k) = 1/2\pi \int_{-\frac{1}{2}L}^{\frac{1}{2}L} dr e^{ikr} e^{-Q(r)} \quad (5.28)$$

$$\cong 1/2\pi \int_{-\infty}^{\infty} dr e^{ikr} e^{-Q(r)}. \quad (5.29)$$

In (5.29) we have passed to the bulk limit  $N, L \rightarrow \infty$ , not an approximation.

At this point our expression for  $\bar{n}_k$  is formally the same as Luttinger's [cf. L (52), L (69)]. The difference is that our  $Q$  is different from his. He obtains  $Q$  by evaluating an infinite Toeplitz determinant with the result that [L (70)]

$$Q(r) = \lambda^2/2\pi^2 \int_0^\infty dp \frac{1 - \cos pr}{p} |v(p)|^2. \quad (\text{Luttinger}) \quad (5.30)$$

Our  $Q$ , which is the correct one to use, is obtained by combining (5.15), (5.17), and (5.25), replacing sums by integrals in the usual way, and using the definition (4.11) of  $\varphi(p)$ . The result is

$$Q(r) = \lambda^2/2\pi^2 \int_0^\infty dp \frac{1 - \cos pr}{p} |u(p)|^2, \quad (5.31)$$

where

$$|u(p)|^2 = (2\pi^2/\lambda^2) \{ (1 - (\lambda v(p)/\pi)^2)^{-\frac{1}{2}} - 1 \}. \quad (5.32)$$

It is worth noting that (5.30) agrees with (5.31) to leading order in  $\lambda^2$ .

Since we have not yet specified  $v(p)$ , we may now follow Luttinger's discussion from this point on with the proviso that we use the correct ( $\lambda$  dependent)  $u(p)$  instead of  $v(p)$ . The reader is referred to pages 1159 and 1160 of Luttinger's paper.

There are two main conclusions one can draw. The first is that if we start with a  $\delta$ -function interaction [so that  $v(p)$  and hence  $u(p)$ ] are constants, it can be shown that  $\bar{n}_k = \frac{1}{2}$  for all  $k$ . Such a result is quite unphysical, but it is not unreasonable because the ground-state energy  $W$  (4.13a) diverges when  $v(p) = \text{constant}$  at large  $p$ . Also, the result would be the same if we started with the more physical interaction

$$H' = 1/L \sum_p \{ \rho_1(p) + \rho_2(p) \} \{ \rho_1(-p) + \rho_2(-p) \} v(p)$$

discussed in the Appendix. This is indeed unfortunate, because relativistic field theories usually begin with local ( $\delta$ -function) interactions.

The second conclusion is that if one makes a reasonable assumption about  $v(p)$ , and hence about  $u(p)$  and  $Q(r)$ , one finds that for  $k$  in the vicinity of  $k_F$ ,  $\bar{n}_k$  behaves like

$$\bar{n}_k \sim d - e |k - k_F|^{2\alpha} \sigma(k - k_F), \quad (5.33)$$

where

$$\begin{aligned} \sigma(k) &= 1, & k > 0 \\ &= -1, & k < 0 \end{aligned} \quad (5.34)$$

and  $d, e$ , and  $\alpha$  are certain positive constants. Now in Luttinger's calculation

$$\alpha = \lambda^2/4\pi^2 v(0)^2, \quad (\text{Luttinger}) \quad (5.35)$$

$$[\text{cf. L(75)}], \quad \text{where } v(0) \equiv \lim_{p \rightarrow 0} v(p).$$

If  $2\alpha < 1$ , then the conclusion to be drawn is that although the interaction removes the discontinuity in  $\bar{n}_k$  at the Fermi surface, we are left with a function that has an infinite slope there. There is, so to speak, a residual Fermi surface. In Sec. IV of his paper, Luttinger shows that at least for one example of  $v(p)$  perturbation theory gives the same qualitative result as (5.33) with the same value of  $\alpha$ , (5.35).

If, on the other hand,  $2\alpha > 1$  then there is no infinite derivative at the Fermi surface.  $\bar{n}_k$  is perfectly smooth there (although, technically speaking, it is nonanalytic unless  $2\alpha = \text{odd integer}$ .) In this case virtually all trace of the Fermi surface has been eliminated. But notice that the correct  $\alpha$  to use is obtained by replacing  $v(0)$  by  $u(0) \equiv \lim_{p \rightarrow 0} u(p)$  in (5.35), i.e.,

$$2\alpha = \{1 - [\lambda v(0)/\pi]^2\}^{-\frac{1}{2}} - 1. \quad (5.36)$$

Thus, even subject to the requirement that  $|\lambda v(0)|$  be less than  $\pi$ ,  $2\alpha$  can become as large as one pleases. Yet perturbation theory predicts (5.35) which yields  $2\alpha$  always less than  $\frac{1}{2}$ .

We may conclude that a strong enough interaction can eliminate the Fermi surface, while perturbation theory predicts that is always there.

## VI. DIELECTRIC CONSTANT

Because the response to external fields of wave vector  $q$  only depends on an interaction expression linear in the density operators, we can immediately obtain for the generalized static susceptibility function or *dielectric constant* (response  $\div$  driving force), for any temperature,  $T$

$$\begin{aligned} \chi_\lambda(q, T) &= \chi_0(q, T) \{\sinh \varphi(q) + \cosh \varphi(q)\}^2 \cosh 2\varphi_q \\ &= \chi_0(q, T) \frac{1}{1 + \lambda v(q)/\pi} \end{aligned} \quad (6.1)$$

in terms of the "unperturbed" susceptibility  $\chi_0(q, T)$ . It is also a simple exercise to calculate exactly the time dependent susceptibility in terms of the "unperturbed" quantity.

It is interesting to note that the susceptibility can diverge (which is symptomatic of a phase transformation) only for

$$\lambda v(q) \rightarrow -\pi, \quad (6.2)$$

i.e. only for sufficiently *attractive* interactions and not for repulsive [ $v(q) > 0$ ] interactions.

Recently Ferrell<sup>6</sup> advanced plausible arguments why a one-dimensional metal cannot become superconducting. We can prove this rigorously in the present model. The electron-phonon interaction is

$$H_{e1-ph} = \sum_p g(p) [\rho_1(p) + \rho_2(p)] \cdot [\xi_p + \xi_{-p}^+], \quad (6.3)$$

where  $\xi$  and  $\xi^+$  are the phonon field operators. In the "filled-sea" limit this coupling is bilinear in harmonic-oscillator operators, and therefore the Hamiltonian continues to be exactly diagonalizable. The new normal modes can be calculated and there is found to be no phase transition at any finite temperature.

## APPENDIX

We shall be interested in extending Luttinger's model in two ways. Firstly, we note that the restriction  $\bar{V} = 0$  is really not necessary. Turning back to Eqs. (2.13) *et seq.* we impose periodic boundary conditions  $\Psi(\dots, x_i + L, \dots) = \Psi(\dots, x_i, \dots)$ , and find that

<sup>6</sup> R. A. Ferrell, Phys. Rev. Letters 13, 330 (1964).

$$(q + N\lambda\bar{V}) \text{ and } (k + M\lambda\bar{V}) = 2\pi/L \times \text{integer} \quad (A1)$$

replace the usual condition (2.16), where  $N$  = number of "1" particles and  $M$  = number of "2" particles. However, when  $N, M \rightarrow \infty$  in the field-theoretic limit the problem evidently becomes ill-defined unless  $\bar{V} \equiv 0$ .

A less trivial observation concerns the form of the interaction potential. There is no reason to restrict it to the form  $\propto \rho_1 \rho_2$ , and in fact the more realistic two-body interaction

$$H' = \frac{\lambda}{L} \sum_p v(p) \{\rho_1(-p) + \rho_2(-p)\} \{\rho_1(p) + \rho_2(p)\} \quad (A2)$$

is fully as soluble as the one assumed in the text, for any strength positive  $v(p)$ , and provided only

$$\lambda v(p) > -\frac{1}{2}\pi, \quad (A3)$$

i.e. provided no Fourier component is *too* attractive. The shift in the ground-state energy is now given by

$$W_2 = \sum_{p>0} p \left\{ \left( 1 + \frac{2\lambda v(p)}{\pi} \right)^{\frac{1}{2}} - 1 \right\}. \quad (A4)$$

The plasmon energy is now

$$\epsilon''(p) \equiv |p| (1 + 2\lambda v(p)/\pi)^{\frac{1}{2}} \quad (A5)$$

and for the important case of the Coulomb repulsion,  $v(p) = p^{-2}$ , the plasmons describe a relativistic boson field with mass

$$m^* \equiv (2\lambda/\pi)^{\frac{1}{2}} \quad (A6)$$

and dispersion

$$\epsilon''(p) = (p^2 + m^{*2})^{\frac{1}{2}}. \quad (A7)$$

Here, too,  $d\epsilon''/dp < 1$ .

## ACKNOWLEDGMENTS

We thank D. Jepsen, L. Landovitz, and T. Schultz for interesting discussions during various stages of this work. We especially thank J. Luttinger for performing a detailed review of his own work and ours, and thus corroborating the results of the present article. While reading our manuscript, Professor Luttinger called our attention to Schwinger's relevant, indeed prophetic remarks (cf. Ref. 5), and we thank him for this reference. We wish to thank Dr. A. S. Wightman, for pointing out to us that P. Jordan first discovered the bosons associated with fermion fields (in his attempts to construct a neutrino theory of light), and for a list of references.<sup>7</sup>

<sup>7</sup> P. Jordan, Z. Physik 93, 464 (1935); 98, 759 (1936); 99, 109 (1936); 102, 243 (1936); 105, 114 (1937); 105, 229 (1937). M. Born and N. Nagendra-Nath, Proc. Ind. Acad. Sci. 3, 318 (1936). A. Sokolow, Phys. Z. der Sowj. 12, 148 (1937).