

# 拓扑相变与拓扑场论作业

**DDL:06-16** 假设 15 周正常结课放假

- 1 阅读文献 (二选一) : A four-dimension generation of quantum hall effect[1] 或 Dirac's monopole wihtout strings: classical lagrange theory [2];
- 2 若  $|\psi_\alpha\rangle = g_{\alpha j}|\psi_j\rangle$ ,  $A' = g^{-1}Ag + g^{-1}dg$ ,  $F' = g^{-1}Fg$ , 证明:  $Q_{2n+1}(A', F') - Q_{2n+1}(A, F) = Q_{2n+1}(g^{-1}dg, 0) + d\alpha_{2n-2}$ ,  $Q_{2n+1}(g^{-1}dg, 0) = \omega_{2n+1}(g)$  NJP,D,065011(2011) eq24,25 [3];
- 3 利用 lundell Table 1, steifel manifold, [4] 推导 NJP,12,065010(2010) table3 [3];
- 4 推导 SSH model 论文 eq.2.1 - eq.3.19 [5], 可参考 Su-Schrieffer-Heeger model applied to chains of finite length[6];
- 5 推导文章 Finite Size Effects on Helical Edge States in a Quantum Spin-Hall System, Zhou Bin, Shunqin Shen. [7];
- 6 推导文章公式和结论 Coupled Spin and Valley Physics in Monolayers of MoS<sub>2</sub> and Other Group-VI Dichalcogenides\_Di Xiao. [8]
- 7 推导文章 Topological Zero-Energy Modes in Gapless Commensurate Aubry-Andre'-Harper Models, Ganeshan et al. [9].

## References

- [1] Shou-Cheng Zhang and Jiangping Hu. "A Four-Dimensional Generalization of the Quantum Hall Effect". In: *Science* 294.5543 (2001), pp. 823–828. DOI: [10.1126/science.294.5543.823](https://doi.org/10.1126/science.294.5543.823). eprint: <https://www.science.org/doi/pdf/10.1126/science.294.5543.823>. URL: <https://www.science.org/doi/abs/10.1126/science.294.5543.823>.

- [2] Tai Tsun Wu and Chen Ning Yang. “Dirac’s monopole without strings: Classical Lagrangian theory”. In: *Phys. Rev. D* 14 (2 1976), pp. 437–445. DOI: [10.1103/PhysRevD.14.437](https://doi.org/10.1103/PhysRevD.14.437). URL: <https://link.aps.org/doi/10.1103/PhysRevD.14.437>.
- [3] Shinsei Ryu et al. “Topological insulators and superconductors: tenfold way and dimensional hierarchy”. In: *New Journal of Physics* 12.6 (2010), p. 065010. DOI: [10.1088/1367-2630/12/6/065010](https://doi.org/10.1088/1367-2630/12/6/065010). URL: <https://dx.doi.org/10.1088/1367-2630/12/6/065010>.
- [4] Albert T. Lundell. “Concise tables of James numbers and some homotopy of classical Lie groups and associated homogeneous spaces”. In: *Algebraic Topology Homotopy and Group Cohomology*. Ed. by Jaume Aguadé, Manuel Castellet, and Frederick Ronald Cohen. Berlin, Heidelberg: Springer Berlin Heidelberg, 1992, pp. 250–272. ISBN: 978-3-540-46772-4.
- [5] W. P. Su, J. R. Schrieffer, and A. J. Heeger. “Soliton excitations in polyacetylene”. In: *Phys. Rev. B* 22 (4 1980), pp. 2099–2111. DOI: [10.1103/PhysRevB.22.2099](https://doi.org/10.1103/PhysRevB.22.2099). URL: <https://link.aps.org/doi/10.1103/PhysRevB.22.2099>.
- [6] Fernando L. J. Vos, Daniel P. Aalberts, and Wim van Saarloos. “Su-Schrieffer-Heeger model applied to chains of finite length”. In: *Phys. Rev. B* 53 (22 1996), pp. 14922–14928. DOI: [10.1103/PhysRevB.53.14922](https://doi.org/10.1103/PhysRevB.53.14922). URL: <https://link.aps.org/doi/10.1103/PhysRevB.53.14922>.
- [7] Bin Zhou et al. “Finite Size Effects on Helical Edge States in a Quantum Spin-Hall System”. In: *Phys. Rev. Lett.* 101 (24 2008), p. 246807. DOI: [10.1103/PhysRevLett.101.246807](https://doi.org/10.1103/PhysRevLett.101.246807). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.101.246807>.
- [8] Di Xiao et al. “Coupled Spin and Valley Physics in Monolayers of MoS<sub>2</sub> and Other Group-VI Dichalcogenides”. In: *Phys. Rev. Lett.* 108 (19 2012), p. 196802. DOI: [10.1103/PhysRevLett.108.196802](https://doi.org/10.1103/PhysRevLett.108.196802). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.108.196802>.
- [9] Sriram Ganeshan, Kai Sun, and S. Das Sarma. “Topological Zero-Energy Modes in Gapless Commensurate Aubry-André-Harper Models”. In: *Phys. Rev. Lett.* 110 (18 2013), p. 180403. DOI: [10.1103/PhysRevLett.110.180403](https://doi.org/10.1103/PhysRevLett.110.180403). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.110.180403>.